

Primordial matter density contrast and the size of the very early universe in the Quantum Big Bang theory of the cosmological constant

Budh Ram
Physics Department
New Mexico State University
Las Cruces, New Mexico 88003, USA

and

Prabhu-Umrao Institute of Fundamental Research
A2/214 Janak Puri, New Delhi 110058, India

Abstract

We calculate the amount of primordial matter density contrast and the size of the very early universe in the recent Quantum Big Bang theory [Arxiv:0705.4549[gr-qc](2007)] of the cosmological constant. We obtain $(\delta\rho/\rho)_M = 1.75 \times 10^{-5}$, *without* the introduction of an adjustable free parameter. Harrison-Zel'dovich k -dependence with $A = 64/9\pi^2 = 0.72$ and $n = 1$ in $|\delta_k|^2 = Ak^n$ arises inherently. The size of the universe with which it enters the classical Friedmann-Robertson-Walker (FRW) phase comes out to be 0.2 cm. We conclude that the hypothesis of classical inflation at an early stage of cosmic evolution is **not** needed.

In a recent paper [1] we constructed the Quantum Big Bang theory of the cosmological constant in which its presently observed very small value $\sqrt{\Lambda/3} = 1.00 \times 10^{-61}$ – appears naturally. (Throughout this paper we use units in which $G = c = \hbar = 1$). The Quantum Big Bang theory [1] is constructed by applying the method [2] earlier given by Ram, Ram and Ram to the classical de Sitter line element. In the resulting quantum theory there appear discrete de Sitter zero-point quantum states with zero-point energies $\frac{3}{2}\omega_0, \frac{3}{2}\omega_1, \frac{3}{2}\omega_2, \dots; \omega_0, \omega_1, \omega_2, \dots$ being the frequencies of *independent* oscillators.

As a result of the first quantum bang (QB) the zero-point energy $\frac{3}{2}\omega_0$ is reduced to $\frac{3}{2}\omega_1$ and the difference $\frac{3}{2}(\omega_0 - \omega_1) = 2(1)\omega_1$ is transformed into one pair of mass quanta, each quantum of frequency ω_1 . As a result of the 2nd QB, the zero-point energy $\frac{3}{2}\omega_1$ is further reduced to $\frac{3}{2}\omega_2$, the difference $\frac{3}{2}(\omega_1 - \omega_2) = 2(2)\omega_2$ being converted to 2 pairs of mass quanta, each mass quantum of frequency ω_2 . And so on, till as a result of the last 43-rd QB, 43 pairs of mass quanta, each of frequency ω_{43} are created.

Everyone of these mass quanta contributes to the matter density ρ_M , the contribution of each quantum being proportional to the *square* of its frequency ω (see [1]).

First we wish to calculate the matter density contrast $(\delta\rho/\rho)_M$ given in terms of the frequency contrasts $\frac{\Delta\omega_n}{\omega_n}$ ($n = 1, 2, \dots, 43$) by the relation

$$\left(\frac{\delta\rho}{\rho}\right)_M = \sum_{n=1}^{43} 2 \left(\frac{\Delta\omega_n}{\omega_n}\right) \quad (1)$$

since the frequencies $\omega_1, \omega_2, \dots, \omega_{43}$ contribute *independently* to the matter density contrast.

In order to do that, let us look closely at the conceptually mathematical nature of a QB.

After the application in [1] of the method [2] to the de Sitter line element the resulting quantum equation ($\omega = \sqrt{\Lambda/3}$)

$$\left(-\frac{1}{2}\frac{d^2}{dr^2} + \frac{1}{2}\omega^2 r^2\right)U = \frac{1}{8\pi}U \quad (2)$$

has only one eigenvalue $\epsilon = \frac{1}{8\pi}$ in

$$\epsilon = \left(2n + \frac{3}{2}\right)\omega, \quad n = 0, 1, 2, \dots \quad (3)$$

Assigning $\epsilon = \frac{1}{8\pi}$ to the $n = 0$ ground state results in an oscillator of frequency $\omega_0 = \frac{1}{12\pi}$. On the other hand, assigning this eigenvalue $\frac{1}{8\pi}$ to the $n = 1$ state results in an oscillator of a different frequency $\omega_1 = \frac{1}{28\pi}$, the relation between ω_0 and ω_1 being

$$\frac{3}{2}\omega_0 = \left(2(1) + \frac{3}{2}\right)\omega_1. \quad (4)$$

Eq.(4) is interpreted as: the first QB transforms the energy difference $\frac{3}{2}(\omega_0 - \omega_1) = 2(1)\omega_1$ to one pair of mass quanta, each mass quantum of frequency ω_1 . In quantum language, the first QB is transition of the isotropic oscillator of frequency ω_1 , from its $|n = 1\rangle$ state to its $|n = 0\rangle$ ground state and spontaneous emission (a la Einstein) of one *p-quantum* (paired-quantum) of frequency ω_1 . This is equivalent to a *Planck* oscillator spontaneously emitting one quantum of frequency ω_1 , and is thus analogous to an atomic transition from one quantum state to another with spontaneous emission of a light quantum of frequency ω_1 (see [3] and references cited therein).

Hence one can calculate the spread $\Delta\omega$ in the frequency ω of the spontaneously emitted mass-quantum in the same manner one calculates the spread in the frequency of the light-quantum in the atomic case, namely, by using the famous *Golden Rule* of time-dependent perturbation theory [4]. Thus

$$\Delta\omega = \omega_{AB} = \int 2\pi\rho_\omega |\langle B|H_I|A\rangle|^2 d\Omega. \quad (5)$$

In Eq.(5), ω_{AB} is the total transition probability per unit time for spontaneous emission of a mass quantum of frequency ω for the isotropic oscillator transition $|A\rangle \rightarrow |B\rangle$,

$$\rho_\omega = \frac{\omega^2}{(2\pi)^3} \quad (6)$$

is the density of states in the ω continuum with the emitted quantum of momentum $\vec{k}$ ($k = \omega$) directed within the solid angle $d\Omega$, and H_I is the self-interaction of the oscillator which we take to be

$$H_I = p_r = \frac{dr}{d\tau} \quad (\tau \text{ is the proper time, see [1]}) \quad (7)$$

analogous to the atomic case.

For the interaction (7), the squared matrix element

$$|\langle B|H_I|A\rangle|^2 = \omega^2 |\langle B|r|A\rangle|^2. \quad (8)$$

Then the contribution $\left(\frac{\delta\rho}{\rho}\right)_n$ from the n -th QB to the total matter density contrast is simply given by

$$\left(\frac{\delta\rho}{\rho}\right)_n = \left(\frac{2\Delta\omega_n}{\omega_n}\right) = \frac{2\omega_n^3}{\pi} |\langle 0|r|n\rangle|^2. \quad (9)$$

For a given n the matrix element $|\langle 0|r|n\rangle|^2$ can be easily calculated using the isotropic oscillator wave functions given in [5] (we explicitly reproduce the first three wave functions in the Appendix). For the first QB

$$|\langle 0|r|1\rangle|^2 = \frac{2}{3\pi\omega_1} \quad (10)$$

giving

$$\left(\frac{\delta\rho}{\rho}\right)_1 = \frac{4\omega_1^2}{3\pi^2} = 1.75 \times 10^{-5} \quad (11)$$

with $\omega_1 = \frac{1}{28\pi}$ (see [1]).

For the second QB

$$|\langle 0|r|2\rangle|^2 = \frac{1}{30\pi\omega_2} \quad (12)$$

giving

$$\left(\frac{\delta\rho}{\rho}\right)_2 = \frac{\omega_2^2}{15\pi^2} = 7 \times 10^{-8} \quad (13)$$

with $\omega_2 = 3.1 \times 10^{-3}$ (see [1]).

Contribution from the 3rd QB is of order 10^{-11} , i.e., negligible, and it is expected to be negligible for the remainder of the QB's in view of how the ω_n 's decrease with increasing n . Thus the major contribution to the density contrast comes from the first QB and $(\delta\rho/\rho)_M$ remains almost constant at the value $\simeq 1.75 \times 10^{-5}$ during the successive 41 QB's. And it enters the classical FRW phase with this value.

With $\omega_1 = k_1 \rightarrow k$, Eq.(11) is rewritten as

$$\left(\frac{\delta\rho}{\rho}\right)^2 = \frac{16}{9\pi^4} k^3 k. \quad (14)$$

Note the Harrison-Zel'dovich k -dependence in (14). Equating (14) with Eq. (7.80) of [6] (see also [7]), namely

$$\left\langle \left(\frac{\delta M}{M}\right)_R^2 \right\rangle = \frac{A}{2} \left(\frac{k^3 k}{2\pi^2}\right)_{k=R^{-1}}, \quad (15)$$

one gets for the Zel'dovich amplitude A in $|\delta_k|^2 = Ak^n$ ($n = 1$) the value [8]

$$A = \frac{64}{9\pi^2} = 0.72. \quad (16)$$

In essence the matter density contrast $\left(\frac{\delta\rho}{\rho}\right)_M \simeq 1.75 \times 10^{-5}$, with its Harrison-Zel'dovich k -dependence [9] and $A = 0.72$, results from the Quantum Big Bang. And then the classical FRW description takes over.

Next we answer the question: what is the size of the universe when it enters the classical FRW phase?

The universe enters the classical FRW description after the **last** 43-rd QB with zero-point energy $\frac{3}{2}\omega_{43} = 1.5 \times 10^{-61}$ (see [1]). The state of the universe with this zero-point energy is described by the radial quantum equation [1]

$$\left[-\frac{1}{2r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{2} \omega_{43}^2 r^2 \right] \psi_{43} = \left(2(43) + \frac{3}{2} \right) \omega_{43} \psi_{43} \quad (17)$$

with a completely spherically symmetric state function ψ_{43} . Thus the size of the universe with which it enters the classical FRW phase, a_{ECP} , is simply $\sqrt{4\pi}$ times [1,10] the square root of the expectation value of r^2 for the state ψ_{43} , viz.,

$$a_{ECP} = \sqrt{4\pi\langle r^2 \rangle_{43}}. \quad (18)$$

The expectation value of r^2 for the state ψ_{43} is most easily calculable by using the well-known result [11] that for a given quantum state of the isotropic oscillator the expectation value of the potential energy equals half the total energy:

$$\left\langle \frac{1}{2}\omega_{43}^2 r^2 \right\rangle_{43} = \frac{1}{2} \left(2(43) + \frac{3}{2} \right) \omega_{43} \quad (19)$$

which, with $\omega_{43} = 10^{-61}$ (see [1]), gives

$$\sqrt{\langle r^2 \rangle_{43}} = 3 \times 10^{31} (= 4.8 \times 10^{-2} \text{ cm}). \quad (20)$$

Or

$$a_{ECP} = 0.2 \text{ cm}. \quad (21)$$

Thus the universe enters the classical FRW phase with a size equal to 0.2 cm.

With the present size of the universe $\simeq 10^{28}$ cm and the present cosmic microwave background (CMB) radiation temperature $\simeq 2.4 \times 10^{-4}$ eV (2.73 K), the temperature and entropy of the very early universe come out to be 10^{16} GeV and 10^{87} , respectively.

It is clear from the results obtained in this paper that the hypothesis of classical inflation [12] at an early stage of cosmic evolution is **not** needed.

Acknowledgments

The author thanks Rajeev Bhalerao, Arun Ram, Nilam Ram, Narayan Banerjee, Subha Majumdar, and Anatoly Klypin for useful conversations, the Tata Institute of Fundamental Research, Mumbai for a pleasant stay, and the Mathematics Department, Melbourne University, Australia for a delightful visit. He is grateful to Rajeev Bhalerao for his continued help and support.

Appendix

Here we reproduce the first three radial wave functions $R_{n,\ell=0}(r)$, $n = 0, 1, 2$, for the isotropic oscillator from [5].

$$\begin{aligned} R_0(r) &= \pi^{-1/4} \alpha^{3/2} 2 e^{-\frac{1}{2}(\alpha r)^2}, \quad \alpha = \omega^{1/2}, \\ R_1(r) &= \pi^{-1/4} \alpha^{3/2} \frac{2^{3/2}}{3^{1/2}} \left(\frac{3}{2} - (\alpha r)^2 \right) e^{-\frac{1}{2}(\alpha r)^2}, \\ R_2(r) &= \pi^{-1/4} \alpha^{3/2} \frac{2^{3/2}}{(15)^{1/2}} \left[\frac{15}{4} - 5(\alpha r)^2 + (\alpha r)^4 \right] e^{-\frac{1}{2}(\alpha r)^2}. \end{aligned}$$

References and Notes

- [1] Budh Ram, ArXiv:0705.4549 [gr-qc] (2007).
- [2] B. Ram, A. Ram, and N. Ram, ArXiv:gr-qc/0504030 (2005); B. Ram, Phys. Lett. A265, 1 (2000). See also, B. Ram and J. Shirley, ArXiv:gr-qc/0604074 (2006).
- [3] B. Ram, ArXiv:quant-ph/0503109 (2005).
- [4] W. Heitler, *The Quantum Theory of Radiation*, third edition (Dover Publications, New York 1984), §17 and 18.
- [5] M.A. Preston and R.K. Bhaduri, *Structure of the Nucleus* (Addison-Wesley Publishing Co. 1975), p.120.
- [6] J.V. Narlikar, *An Introduction to Cosmology*, 3rd edition (Cambridge Univ. Press 2002), p.258.
- [7] T. Padmanabhan, *Structure Formation in the Universe* (Cambridge Univ. Press 1993), Chap.5.

- [8] The best fit for the power-law LCDM model for WMAP + 2d FGRS + $\text{Ly}\alpha$ data gives $A \simeq 0.75$; see D.N. Spergel *et al.*, Ap J. Supplement Series 148, 175 (2003).
- [9] The Harrison-Zel'dovich model is consistent with the WMAP three-year data; see W.H. Kinney, E.W. Kolb, A. Melchiorri, and A. Riotto, ArXiv:astro-ph/0605338 (2006).
- [10] H.P. Robertson, Phil. Mag. **5**, 835 (1928).
- [11] L.I. Schiff, *Quantum Mechanics*, 3rd edition (McGraw-Hill, New York, 1955), p.180.
- [12] A.H. Guth, Phys. Rev. **D23**, 347 (1981).